

**Eonlipid**®

FOOD SUPPLEMENT AGAINST LIPID AND METABOLIC DISORDERS, OXIDATIVE PROCESSES AND FREE RADICALS





EONLIPID is a multicomponent food supplement based on Hydroxytyrosol, Maqui, Amla, Monacolin K and herbal extracts.

## WHEN USING EONLIPID

On the basis of the international literature it is indicated:

- in cardiovascular and brain degenerative diseases
- for the activation of **lipid / glycidic metabolism** and against metabolic syndrome
- against the processes of **atherogenesis**
- to combat **inflammaging** (senescence and cellular inflammation)
- against **oxidative stress** and **free radicals**



## THE NATURAL INGREDIENTS OF EONLIPID®



Hydroxytyrosol is derived from olive and it has proven antioxidant, antidyslipidemic (1) properties

and contributes to regulate blood pressure and circulation (2); hydroxytyrosol is an innovative natural ingredient extremely promising in metabolic syndrome (3) and in the management of "inflammaging" (inflammation causing aging process).

Maqui (Aristotelia chilensis, Mol.Stuntz) is an



energizing berry with multiple beneficial effects on metabolism (4), and it is probably one of the most

powerful natural principles in terms of antioxidant power (5) and thus it is very useful in the fight against free radicals (6).

## Amla

(Phyllanthus Emblica) forms the basis of nutritional therapy in Ayurvedic



medicine, by virtue of its certified antioxidant power (7) and its efficacy in hyperdislipidemias (8) and in metabolic syndrome (9).

Monacolin K. extracted from red yeast rice (Monascus purpureus, Went), is a natural statin with a documented lipid (blood cholesterol namely) lowering effect (10).



Berberine is another natural substance with proven blood lipid lowering effect (11), while

## Coenzyme Q10 (ubiquinol)

Serum Antioxidant Capacity

2246.06

S.D. 192.8

+6.8%

0 day

p = 0.0042

2245

2220

2195

2170

2145

2120

2095

2070

provides special protection to heart and muscle functions (12). The administration of Folic Acid contributes to regulate methylation processes (which may damage DNA) and the normal homocysteine metabolism, which interacts with athero-thrombotic processes (13).

Astaxanthin with its high antioxidant power (14) complete the list of the natural principles of Eonlipid.


## Observational PILOT STUDY

A polyphenol-based multi-component nutraceutical in dysmetabolism and oxidative stress; results from a pilot study, Corsi R, Mosti G; Cavezzi A; Urso S.U; Dimitrova G; Fioroni; Colucci R; Quinzi V J of Dietary Supplements 2017

310

### **Total Cholesterol** 250 -20° 240 230 220 210 200 194.83 190 180 170 60 days





0 day p < 0.0001

60 days

110

100

## Oxidized Cholesterol



p = 0.0511

# Free Radicals



p = 0.036

## Glycemia



p = 0.0352

EONLIPID is a unique supplement: 8 natural ingredients in a synergic anti-aging action and to combat lipid profile disorders and metabolic alterations.



Maqui berry (Aristotelia chilensis [Mol] Stuntz) freeze-dried fruit; olive involution private and incisor in more state in the recent and incisor tree (Olea europaea L.) leaves d.e. (10-12% hydroxytyrosol). Charging agent: microcrystalline cellulose: Barberry (Berberis vulgaris L.) bark de. (98% berberine); Amal (Phyllanthus emblica L.) fruit de. (30% of tannins); Ubiquinol (coenzyme O10 reduced); fermented red rice (Monascus purpureus Went) microfunghi d.e. (5% monakoline K); Haematococcus (Haematococcus pluvindis Flotow) microalgae d.e. (2.5% astaxanthin); folic acid; anti-caking agents: vegetable magnesium sterates, colloidal silica. Dr. vetratest are supported on e, colloidal silica. Dry extracts are supported or corn maltodextrins.

Serving size: 1 or 2 tablets a day between meals.

## Warnings:

Warnings: Warnings: Warnings: do not consume a daily quantity equal to or greater than 3 mg of monacolins from red fermented rice [1 tablet of Eonlipid contains 1 mg of monacolin K). It should not be consumed by pregnant or breastfeeding women, children under the age of 18 and adults over the age of 70. It should not be consumed if you are taking medicines to large extensions and the state of the second of the state of the second of to lower cholesterol or if you already consume other products conta on long et ploase trie. Consult a doctor about consuming this product if you experience any health problems. Flood supplements should not be understood as substitutes for a varied, balanced diet and a healthy lifestyle. Do not exceed the recommended daily doses. Keep out of reach of children under 3 years of age.

## Content of active ingredients

60 days

<u> </u>		
Ingredients	1 tablet	2 tablets
olive leaves dry extract	75 mg	150 mg
corresponding to Hydroxytyrosol	7.5 mg	15 mg
maqui dry extract	150 mg	300 mg
amla fruit dry extract	100 mg	200 mg
corresponding to tannins	30 mg	60 mg
red yeast rice dry extract	20 mg	40 mg
corresponding to monacolin K	1 mg	2 mg
barberry root bark dry extract	125 mg	250 mg
corresponding to berberine	122.5 mg	245 mg
haematococcus algae dry extract	10 mg	20 mg
corresponding to astaxanthin	0.25 mg	0.5 mg
coenzyme Q10 (ubiquinol)	50 mg	100 mg
folic acid	100 mcg	200 mcg

EONLIPID supplement tablets are packaged in blisters. One pack contains 30 tablets. EONLIPID does NOT contain gluten or sweeteners, it is GMO FREE and Vegan Ök

Produced in via Agrate, 21 - Concorezzo (MB) - Packaged in v. Maniscalchi 5 - Carpi (MO)







## BIBLIOGRAPHY-

1) Can red yeast rice and olive extract improve lipid profile and cardiovascular risk in The art of the vest in the art of the whole the controlled randomized the late Vehicles of the Vehicles of the

terarcuppeanbounnavirunizung zu park 1: F12
3) Beneficial effects of the olive oil phenolic components oleuropein and hydroxytyrosol: focus on protection against cardiovascular and metabolic diseases Bullotta S, Celano M, Lepore SM et al Transi Med. 2014 Aug 3;72:719
4) Nutraceutical and antioxidant effects of a delphinidin-rich maqui berry extract Delphinol: a review. Watson RR, Schönlau F Minerva Cardioangiologica 2015, 63(2) Suppl:1:1-12

Jouppil 172 5) Phytochemical profile and the antioxidant activity of Chilean wild black Berry fruits, Aristotelia chilensis (Mol) Stunz (Elaeocarpaceae). Cespedes C, Valdes-Morales M, Avila JG et al Food Chemistry 2011,

6) A Randomized Clinical Trial Evaluating the Efficacy of an Anthocyanin-Maqui Berry

Extract (Delphino!) on Oxidative Stress Biomarkers Davinelli S, Bertoglio JD, Zarrelli A et al, Journal of the American College of Nutrition, 2015; M. Ox 1, 28–33 '7) The total antioxidant content of more than 3100 foods, beverages, spices, herbs and supplements used worldwide Carlsen MH, Halvorsen BL, Holte K et al, Nutr J. 2010

8) A comparative clinical study of hypolipidemic efficacy of Amla (Emblica officinalis) o/Acomparative climina study of hypothemic measor of him a climina of him a with 3-hydroxy-3-methylglutaryl-coenzyme-A reductase inhibitor simvastatin Gopa B, Bhatt J, Hemavathi KG, Indian J Pharmacol. 2012 Mar-Apr, 44(2): 238–242.

9) Supplementation of a standardized extract from Phyllanthus emblica impro cardiovascular risk factors and platelet aggregation in overweight/class-1 obese adults. Khanna S, Das A, Spieldenner J et al. J Med Food. 2015 Apr;18(4):415-20 aouris, marina aj posa, spierucinire) et al. Preu Prouz, 2013 pij, 16(4):415-20 1D) Effect of red yeast rice combined with antioxidants on lipid pattern, hs-CRP level, and endothelial function in moderately hypercholesterolemic subjects. Cicero Af, Morbini M, Parini Aet al, Ther Clin Risk Manag. 2016 Feb 23;12:281-6. 11) The effects of berberine on blood lipids: a systemic review and meta-analysis of randomized controlled trials. Dong H, Zhao Y, Zhao L et al Planta Med. 2013 Apr,79(6):437-46

Agry79(c) 437-46

12). Coenzyme (101 and Utility in Heart Failure: Just Another Supplement? Oleck S, Ventura HD. Curr Heart Fail Rep. 2016 Aug;13(4):100-95

13). DACH-LIGA homocystein (german, austrian and swiss homocysteine society): consensus paper on the rational clinical use of homocysteine, folic acid and 8-vitamins in cardiovascular and thrombotic diseases: guidelines and recommendations. Stanger 0, Herrmann W, Pietrzik K et al, DACH-LIGA Homocysteine. V. Clin Chem Lab Med. 2003 Nov;41(17):1392-403

14) Free Radioal Scavenging and Cellular Antioxidant Properties of Astaxanthin. Dose L Matsupo S, Vokokawa H et al, Int I Mal Sci. 2016 Ian 14:17(1)

15). A polyhenol-based multi-component nutraecutical in dysmetabolism and oxidative stress: results from a pilot study. Corsi R, Mosti G; Cavezzi A; Urso S.U; Dimitrova G, Horoni; Colucci R, Quinzi V Journal of Dietary Supplements 2017, Accepted for publication