

Eon-DK2®

FOOD SUPPLEMENT BASED ON
VITAMIN D2 AND VITAMIN K2, IN
GRAPESEED OIL, USEFUL FOR
OPTIMIZE THE CALCIUM METABOLISM
AND THE IMMUNE SYSTEM



Eon-DK2®

is a food supplement unique
for formulation, dose and
quality: 5 drops contain 50
mcg of vitamin D3
(equivalent to 2.000 I.U.)
and 50 mcg of vitamin K2

EON-DK2®

Great quality of active ingredients: vitamin D3 (cholecalciferol) of natural origin and vitamin K2 (menaquinone) of the MK-7 type, from natural fermentation.

Adjustable dosage according to needs, thanks to the dropper.

Bioavailability increased by sublingual administration.

Better absorption of vitamin D3 and K2, due to the dilution in grapeseed oil, extracted from the seeds of grape berries.

The integration of vitamin D3 in association with vitamin K2, can contribute to optimize the metabolism of calcium in the ossification process, favoring the deposition mainly in the bones. The synergy of the two vitamins helps to improve the functions of the immune and cardiovascular systems



WHEN TO USE EON-DK2

INDICATIONS OF USE, ACCORDING TO EVIDENCE BASED MEDICINE:

- improvement of the immune system (3,4)
- reduction of the risk of osteoporosis (1,2), favoring calcium deposition mainly at bone level, and less in soft tissues (5,9)
- contrast of the process of calcification and sclerosis of the arterial wall, helping to prevent the risk of cardiovascular disease (10,12)
- contribution to normal muscle function and maintenance of normal bones

Tenore degli ingredienti attivi

Ingredients	5 drops
Vitamin D3 (cholecalciferol)	50mcg (2,000 I.U.)
Vitamin K2 (Mk-7 menaquinone)	50mcg

Eon-DK2 is produced in Italy, notified to the Ministry of Health; Eon-DK2 is gluten-free, OGM-free, lactose and food preservatives free



Bottle of 30 mL with dropper

Produced and packaged in
via Agrate 21
Concorezzo (MB) - ITALY
Distribution by Proeon S.u.r.l.
www.proeon.it



EON-DK2

is a unique food supplement

- optimal formulation and concentration: 5 sublingual drops per day contain 50 mcg of vitamin D3 (equivalent to 2,000 I.U.) and 50 mcg of vitamin K2

- adjustable dosage according to needs by dropper

- increased bioavailability due to the possibility of sublingual administration (13,14)

Ingredients:

seed oil (linoleic acid 61%, oleic acid 26%); vitamin K2 (menaquinone), type MK-7, from natural fermentation; vitamin D3 (cholecalciferol); aroma.

Serving size:

5 drops per day orally, preferably sublingually, during meals.

Warnings:

Food supplements are not substitutes for a varied and balanced diet and a healthy lifestyle. Do not exceed the recommended daily dose.

It is recommended to consult your doctor in case of maternity, lactation or for different dosages according to the actual need for vitamin D deficiency. Keep out of reach of children under 3 years of age. Store in a cool, dry place, away from light.

BIBLIOGRAPHY

1) Vitamin D Deficiency. Sizar O, Khare S, Goyal A, Givler A. Vitamin D Deficiency. 2022 Jul 18. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan.
2) Vitamin D and health - The missing vitamin in humans. Chang SW, Lee HJ. Pediatr Neonatol. 2019 Jun.
3) Evidence Regarding Vitamin D and Risk of COVID-19 and Its Severity. Mercola J, Grant WB, Wagner CL. Nutrients. 2020 Oct 31.
4) Vitamin D3 and K2 and their potential contribution to reducing the COVID-19 mortality rate. Goddek S. Int J Infect Dis. 2020 Oct.
5) Vitamin K status and vascular calcification: evidence from observational and clinical studies. Shea MK, Holden RM. Adv Nutr. 2012 Mar 1.
6) Dietary Intake of Menaquinone Is Associated with a Reduced Risk of Coronary Heart Disease: The Rotterdam Study.

Geleijnse JM, Vermeer C, Grobbee DE, Schurgers LJ, Knapen MH, van der Meer IM, Hofman A, Witteman JC. J Nutr. 2004 Nov.
7) A high menaquinone intake reduces the incidence of coronary heart disease. Gast GC, de Roos NM, Sluijs I, Bots ML, Beulens JW, Geleijnse JM, Witteman JC, Grobbee DE, Peeters PH, van der Schoor YT. Nutr Metab Cardiovasc Dis. 2009 Sep.
8) High dietary menaquinone intake is associated with reduced coronary calcification. Beulens JW, Bots ML, Atsma F, Bartelink ML, Prokop M, Geleijnse JM, Witteman JC, Grobbee DE, van der Schoor YT. Atherosclerosis. 2009 Apr.
9) Menaquinone-7 supplementation improves arterial stiffness in healthy postmenopausal women: A double-blind randomised clinical trial. Knapen MH, Braam LA, Drummen NE, Bekers Q, Hoeks AP, Vermeer C. Thromb Haemost. 2015 May.
10) The combination effect of vitamin K and vitamin D on human bone quality: a meta-analysis of randomized controlled trials.

Kuang X, Liu C, Guo X, Li K, Deng Q, Li D. Food Funct. 2020 Apr 30.
11) The Synergistic Interplay between Vitamins D and K for Bone and Cardiovascular Health: A Narrative Review. Van Ballegooijen AJ, Pilz S, Tomaschitz A, Grubler MR, Verheyen N. Int J Endocrinol. 2017
12) Role of vitamin K2 in bone metabolism: a point of view and a short reappraisal of the literature. Capozzi A, Scambra G, Migliaccio S, Lello S. Gynecol Endocrinol. 2020 Apr.
13) A randomized two way cross over study for comparison of absorption of vitamin D3 buccal spray and soft gelatin capsule formulation in healthy subjects and in patients with intestinal malabsorption. Satia MC, Mukim AG, Tibrewala KD, Bhavsar MS. Nutr J. 2015 Oct 29.
14) Rate of change of circulating 25 hydroxyvitamin D following sublingual and capsular vitamin D preparations. Williams CE, Williams EA, Corfe BM. Eur J Clin Nutr. 2019 Dec.